

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 5/10, 15/16, 15/80	A1	(11) International Publication Number: WO 93/25663 (43) International Publication Date: 23 December 1993 (23.12.93)
(21) International Application Number: PCT/US93/05866 (22) International Filing Date: 17 June 1993 (17.06.93) (30) Priority data: 07/899,689 17 June 1992 (17.06.92) US (71) Applicant: UNIVERSITY OF HAWAII [US/US]; Department of Genetics, BioMed A 103, Biomedical Sciences Building, 1960 East-West Road, Honolulu, HI 96822 (US). (72) Inventors: STUART, W., Dorsey ; 1960 East-West Road, Honolulu, HI 96822 (US). KOO, Kenneth ; 18424 Keswick Street, #6, Reseda, CA 91335 (US). MANDEL, Morton ; 2937 Kalakaua, Apt. 27, Honolulu, HI 96815 (US). KATO, Elie, K. ; 2585 Dole Street, Apt. B 214, Honolulu, HI 96822 (US). GREENWOOD, Fred, C. ; 3498 Alani, Honolulu, HI 96822 (US).		(74) Agents: MURASHIGE, Kate, H. et al.; Morrison & Foerster, 2000 Pennsylvania Avenue, N.W., Washington, DC 20006 (US). (81) Designated States: AT, AU, BB, BG, BR, CA, CH, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, PL, RO, RU, SD, SE, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: USE OF <i>mtr</i> GENE SEQUENCES FOR EXPRESSION OF FOREIGN GENES (57) Abstract Expression systems useful in recombinant hosts are provided which permit selection of homokaryotic recombinants capable of producing a desired foreign gene. The expression system contains its own selectable marker. A first nucleotide sequence encoding the desired foreign gene is included in an expression system containing a second nucleotide sequence derived in an illustrative embodiment from the <i>mtr</i> gene of <i>N. crassa</i> which is disposed with respect to the first sequence so as to effect homologous recombination of both the first and second sequences into the <i>mtr</i> locus of the host cell, thus disrupting the endogenous locus. Since the functional product of the <i>mtr</i> gene is required in order to make the cells susceptible to certain poisonous substances, homokaryotic recombinants are resistant to these substances. Successful recombinants can therefore be efficiently selected. The inserted expression system is capable of effecting the production of the desired protein.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

USE OF mtr GENE SEQUENCES FOR
EXPRESSION OF FOREIGN GENES

This invention relates to the use of cloned
5 sequences of the mtr gene of Neurospora crassa to
express DNA encoding heterologous proteins, and to use
of recessive traits to assure homokaryotic
transformation in polynucleated cells.

10 BACKGROUND OF THE INVENTION

The translocation of aliphatic and aromatic
amino acids across the plasma membrane of the
ascomycete, Neurospora crassa, requires a functional
gene product of the genomic mtr locus. Perkins, D.D.
15 et al., Microbiol Rev (1982) 46:426-570. Mutations at
this locus result in defective transport of neutral
aliphatic and aromatic amino acids and the fungus
exhibits altered cell-surface glycoproteins. These
mutations can be selected by resistance to
20 4-methyltryptophan (4-MT) and p-fluorophenylalanine
(pFPA). These substances are poisons, which cannot be
transported into the cell in the absence of the mtr
gene product.

The mtr locus has been mapped to the right
25 arm of linkage group IV by Stadler, H., Genetics
(1966) 54:677-685. Intralocus recombination of
allelic mutants in the mtr locus has been shown by
Larimer, F.W. and DeBusk, A.G., J Bacteriol (1977)
129:1636-1638.

30 A restriction map of a 2.9 kb fragment of
genomic DNA encompassing the mtr coding region has
been produced from a cosmid library of genomic DNA of
Neurospora crassa by Stuart, W.D., et al., Genome
(1988) 30:198-203. In a later paper, Koo, K. and
35 Stuart, W.D., Genome (1991) 34:644-651, the complete
nucleotide sequence of the open reading frame and the
upstream and downstream regulatory sequences of the
mtr locus of an N. crassa strain were described, along
with the location of useful restriction sites. The

disclosure of these articles is incorporated herein by reference.

The 1991 article contains two minor errors in the DNA sequence there provided. They are in the upstream region of what is reported as the open reading frame. Correction of these errors results in extension of the open reading frame further into the 5' region; however, this change does not materially affect the utilization of this gene in the invention described herein. The disclosure of the parent application herein, U.S. serial no. 07/899,689 filed 17 June 1992 is also incorporated herein by reference.

Other transport related genes that confer susceptibility to certain amino acid analogs in N. crassa are also known, such as the pmb locus, which confers canavanine susceptibility.

The present invention provides materials and methods to take advantage of the necessity for a functional mtr gene product to confer susceptibility to certain metabolic poisons such as 4-MT and pFPA. By using as host a cell which is capable of providing functional mtr gene products, successful homologous recombination of an expression system for a desired protein into all functional mtr loci present in the host can be selected for directly.

DISCLOSURE OF THE INVENTION

The invention provides materials and methods for efficient transformation and selection of hosts that provide functional mtr or other recessive gene products, especially filamentous fungal hosts, which are modified to contain an expression system for a desired recombinant foreign protein. Expression systems are used for transformation which are simultaneously capable of producing the desired protein and providing a selectable marker for successful homokaryotic transformants. The expression systems are designed so as to effect homologous

recombination into native mtr loci or analogous loci in the host.

Thus, in one aspect, the invention is directed to an expression system for a gene encoding a desired protein. The expression system consists of a DNA molecule which comprises a first nucleotide sequence encoding the desired protein and a second nucleotide sequence derived from a gene that confers a recessive susceptibility to a poison, such as the mtr gene of N. crassa. The first and second sequences are disposed in relationship so as to permit homologous recombination of the first and second sequences into the target locus of the host in such a way as to incapacitate normal expression of the locus. This prevents the production of functional protein from that locus. The first, coding, nucleotide sequence is operably linked to control sequences which effect expression of the coding sequence when contained in a recombinant host cell. When all functional native, recessive loci are destroyed, the transformed host can be selected by growth on a medium containing a substrate which is poisonous to the cell requiring the gene product for transport. This general method for selecting homokaryotic transformed cells is illustrated herein using the mtr locus of N. crassa.

In other aspects, the invention is directed to filamentous fungal cells modified to contain the expression system of the invention, to methods to obtain such modified cells, and to methods to produce the desired protein using the modified cells.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide sequence of an approximately 2.9 kb fragment of the N. crassa mtr gene containing the entire open reading frame as well as the promoter and transcription terminating signals.

Figure 2 shows nucleotide and deduced amino acid sequences of the open reading frame of the mtr locus.

MODES OF CARRYING OUT THE INVENTION

The present invention provides, in a preferred embodiment, the relevant DNA of a wild type Neurospora crassa mtr gene locus. Figure 1 shows the complete nucleotide sequence of this wild type mtr; an approximately 2.9 kb segment of DNA derived from a cosmid library of N. crassa FGSC No. 3043 is depicted. Because of considerable homology among various strains of N. crassa and across various filamentous fungi, both allelic variants of the nucleotide sequence in Figure 1 and variants derived from alternative strains of filamentous fungal species and other organisms which exhibit unicellular growth may also be used in the invention. Provision of the sequence information of Figure 1 is adequate to permit retrieval of these alternative mtr genes. Modifications of the depicted sequence can be made by site-directed mutagenesis and portions or all of this or other retrieved sequences may be synthesized using well known solid phase nucleotide synthesis techniques. Alternate sequences can also be derived from cosmid libraries prepared from other fungi in a manner analogous to that described for the retrieval of the gene disclosed herein by Koo, K. et al., Genome (1991) 34:644-651, cited above.

Thus, as used herein, the phrase "a nucleotide sequence derived from the mtr gene of N. crassa" refers to a nucleotide sequence as shown in Figure 1, or a functional portion thereof, or a modified form thereof which contains sufficient homology to the sequence depicted to be functional in effecting homologous recombination with a native mtr locus, allelic variants thereof, corresponding loci from other organisms which are retrievable using the depicted sequence as a probe or using relevant portions of the sequence shown in Figure 1 as primers for PCR cloning. Functional portions of all of the foregoing may be used. These nucleotide sequences may be obtained by cloning the retrieved genes, modifying

retrieved genes and amplifying them, or partially or completely synthesized using standard polynucleotide synthesis techniques.

The expression systems of the present invention contain, in addition to a nucleotide sequence derived from the mtr gene of N. crassa, a gene encoding a desired protein to be produced recombinantly in a host cell. Any desired protein can be produced using the expression systems of the invention. Such proteins include both plant and animal proteins and proteins derived from microorganisms. Thus, genes encoding enzymes such as thrombin, tissue plasminogen activator, streptokinase, urokinase, alcohol dehydrogenase, various proteases such as trypsin, chymosin, enterokinase, and the like are suitable for production using the expression systems of the invention. Such enzymes may be useful as pharmaceuticals, as reagents, or as industrial enzymes. Other exemplary types of proteins whose production may be desired include growth factors and cytokines such as interleukin-2, bovine growth hormone, human growth hormone, fibroblast growth factor, TGF- α , TGF- β , bone morphogenic protein, IL-3, IL-6, and the like. In addition, the expression system may be used for the recombinant production of antibodies or fragments thereof, including chimeric antibodies as well as hormones such as insulin, ACTH, the gonadotropins, and the like. Any protein for which a practical means of production is desired and the coding sequence for which is available can be employed in the expression systems and methods of the invention.

The host cells which are useful in the invention method for production of recombinant protein using the expression systems containing nucleotide sequences derived from the mtr gene of N. crassa are cells containing genes encoding functional mtr products which confer susceptibility on the native host to substances toxic to the cells when internalized using the mtr system, such as the 4-MT

and pFPA described above. The host cells can be derived from any species, provided the relevant gene is present in these cells, and provided the cells can be grown in culture. Filamentous fungal host cells
5 are preferred. N. crassa host cells are particularly preferred.

Cells subject to the general method described herein (selection for homokaryotic recombinants by effecting destruction of an endogenous
10 recessive gene conferring susceptibility to a poison) may be any cells which contain such recessive genes. In a manner analogous to that described herein using a nucleotide sequence derived from the mtr locus, sequences derived from the relevant recessive gene
15 locus are used to effect homologous recombination of a desired DNA into the genome of the host cell. The recombination results in destruction of the functionality of the native gene and thus converts the successful homokaryotic transformant into a cell
20 resistant to the poison. Only homokaryotic transformants will therefore grow on media containing the poison.

Other recessive genes of this type, besides the mtr locus, are known. For example, N. crassa also
25 contains a locus encoding a "pmb" product which effects the transport of basic amino acids into the cells. Canavanine, a basic substance which is toxic to native N. crassa, is then not toxic to cells with each such locus destroyed. Successful transformants
30 can then be identified by successful growth on a medium containing canavanine.

The method disclosed herein for efficient selection of successful homokaryotic transformed cells is particularly useful where the cell to be
35 transformed has more than a single copy of a gene locus. Thus, this approach is useful in transformation of diploid or polyploid cells, or, especially, in polynucleated cells such as those found particularly in filamentous fungal species. Since all

of the native loci must be destroyed in order to render the cells resistant, the selection method automatically provides homokaryotic transformants.

Unless otherwise stated, the present invention employs molecular biology, microbiology and recombinant DNA techniques as described, for example, Maniatis, Fritsch & Sambrook, "Molecular Cloning: A Laboratory Manual" (1982); "DNA Cloning: A Practical Approach," Volumes I and II (D.N. Glover ed. 1985); "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization" (B.D. Hames & S.J. Higgins eds. 1985); "Transcription And Translation" (B.D. Hames & S.J. Higgins eds. 1984); "Animal Cell Culture" (R.I. Freshney ed. 1986); "Immobilized Cells And Enzymes" (IRL Press, 1986); B. Perbal, "A Practical Guide To Molecular Cloning" (1984).

The expression systems of the invention provide for insertion of heterologous DNA into cells (including spheroplasts or protoplasts) that can be multinuclear, so that a selection of homokaryotic transformants is facilitated. Homokaryotic transformants may be selected in one step from a colony or mixture containing multinuclear cells. This is particularly advantageous in dealing with Neurospora or other filamentous fungi having vegetative spores (conidia) which contain one or more nuclei. The transformation of such spores, after digestion of the cell walls to produce spheroplasts, is a random event. If the spheroplasts are multinucleated, the transforming DNA may be inserted into less than all of the nuclei that are present. Other multinucleate cells can also be used as subjects for the invention.


Transformation selection protocols directed to a dominant phenotype, such as the restoration of an enzyme which confers the ability to grow on minimal medium, do not select for homokaryotic transformants. Transformation of any single nucleus in any spheroplast will result in growth of transformed

colonies on selective media, even though many of the transformants will also carry untransformed nuclei in a number equal to, or in excess of, the desired transformed nucleus. In the invention method, when a
5 recessive allele is inserted, the recessive genotype can be expressed phenotypically (and thus selected) only if it is the only allele in the culture of transformed strains, thus ensuring that selected transformants will carry only the nuclei which have
10 the constructed gene of interest.

The illustrative expression system herein provides such a recessive allele. Neurospora cells that include wild-type mtr cannot grow on media containing 4-MT or pFPA because the transport system
15 in the host of which the mtr gene product is a part brings these substances into the cell and the cell dies. If the host cells are made homokaryotic mtr negative mutants, the transport system is inoperative so the cell can survive on the media containing these
20 substances, because it is not transported into the cell.

For colonies of multinucleated spheroplasts, transformation of one nucleus to mtr⁻ by insertion of a vector according to the present invention would
25 still result in selection against that transformant because the nontransformed nuclei would continue to produce the mtr protein, the cell will continue to transport the poison and would therefore die as a result. Only if a transformed spheroplast has all mtr
30 loci disrupted will it be able to resist the poison, survive the selection and grow into a strain where all the nuclei contain the inserted gene. Thus, in selecting for resistant transformants, selection of homokaryotes is assured.

35 The general features for use of the expression systems of the invention can be understood in terms of the diagram showing the features of the mtr gene (common features with genes in general).



The 5' untranslated region (5'UT) transcribed into the mRNA contains features which facilitate translation of the mRNA. This is followed by the open reading frame (ORF) which contains the coding region which is ultimately translated into the protein. In the mtr gene illustrated, the ORF also contains a 59-nucleotide intron which remains in the mRNA but is not translated. (See Figure 2.)

Presumably, the intron contributes to enhanced stability of the mRNA. The transcribed region is completed by the 3' untranslated region (3'UT) which contains the polyadenylation signal also contributing to mRNA stability and facilitating translation.

In the DNA, the transcribed region is preceded by the promoter (P) and followed by the transcription terminating sequences (T). The transcription terminating sequences are believed to render expression more efficient, but do not appear to be necessary to effect at least some expression.

The expression systems of the invention contain some or all of the mtr sequence just discussed, or the corresponding portions of other recessive loci, and thus provide for the insertion of DNA comprising the sequence encoding the desired protein to be inserted somewhere in the overall gene or contiguous with one of its termini, provided sufficient deletion or interruption is made in the gene sequences so as to disrupt expression of the wild type gene when homologous recombination occurs.

While homologous recombination can be effected by extension at only one terminus of the desired DNA to be inserted, insertion is more efficient when the DNA to be inserted is bracketed by homologous sequences. The efficiency of homologous recombination is increased with increasing length of nucleotide sequence bearing homology to the target for recombination. It is believed that a minimum of ten

nucleotides is required, but preferably 20, more preferably 100, still more preferably 200 and still more preferably 1000 nucleotides can be included in the construct to effect recombination. The length of the nucleotide sequence provided is arbitrary; the frequency of recombination is simply increased as the length of homologous sequence increases. Furthermore, the homologous sequences, such as the mtr sequences, must be modified in some way so as to make them nonfunctional if they are to be simply ligated to one end of the inserted sequence, rather than automatically disrupted by the DNA to be inserted. Thus, expression vectors which result from the insertion into the homologous sequence of the DNA comprising the gene encoding the desired protein are preferred.

It is, of course, not necessary to utilize the complete mtr gene sequences illustrated; insertion can be made into the mtr gene in which deletions have been made. For example, substantial portions of the open reading frame of the mtr gene may be removed prior to (or after) insertion of the desired DNA.

Using the above diagram for mtr as an illustration, the position for insertion of the DNA comprising the desired coding DNA is arbitrary, but will depend on the nature of the DNA to be inserted. If the inserted DNA consists essentially of the desired coding sequence, to be placed under control of the mtr promoter, the insert must be made so that the coding sequence is in operable linkage with the mtr promoter. Similar comments apply to utilization of the mtr transcription terminator. Thus, the insert is made into the transcribed region of the mtr gene, preferably into the open reading frame. It is possible, but not necessary, to provide for the production of a fusion protein by ligating the desired coding region in open reading frame with the mtr ORF. Alternatively, the DNA comprising the desired coding sequence can be inserted under the control of its own

promoter and/or transcription terminator which, themselves, may either be homologous with the coding sequence or heterologous to it. The promoter must, of course, be functional in the host. An additional copy of the mtr promoter could also, of course, be used. Similarly, DNAs comprising coding sequences for two or more desired heterologous proteins can be inserted, some or all under control of the promoter and/or transcription terminating sequences in the mtr sequence and some or all under control of their own transcription control sequences.

The functional features of the expression system in effecting homologous recombination can be illustrated in the diagrams below which show but two of many options for the construction and function of such vectors:

In the first illustration, the vector contains the transcribed portion of a mammalian gene as a replacement for the transcribed portion of the mtr gene. Upon homologous recombination, the transcribed sequences of the mammalian gene are integrated into allele replacing the corresponding regions of the mtr gene and thus disrupting its function. This is shown diagrammatically below:

vector: ---PPPP-MAMMALIAN GENE-TTTTTT----

integrates into

genome: ---PPPPP-MTR GENE-TTTTTT-----

at the mtr locus disrupting the mtr gene and producing the allele:

---PPPPP-MAMMALIAN GENE-TTTTTT---

35

In a second alternative, multiple genes under control of their own promoters and terminators can be inserted between the controls of the host mtr gene as shown in the following diagram:

vector:---PPPPP-GENet-pGENet-pGENet-pGENet-TTTTT

integrates into

5 genome: ---PPPPP-MTR GENE-TTTTTT-----

at the mtr locus disrupting the mtr gene and producing the allele:

10 ---PPPPP-GENet-pGENet-pGENet-pGENet-TTTTTT---.

In this illustration, only the first gene in the sequence is under control of the mtr promoter; the remaining genes bear their own control sequences.

15 Again, the function of the wild type mtr gene will be destroyed.

Of course, substantial portions of the open reading frame of the mtr gene can also be retained, if desired, so long as insufficient sequences are provided to effect production of a functional protein gene product. Indeed, deletions, additions and alterations in the sequence encoding the mtr protein product may be made in the expression system constructed so long as the remaining sequences fail, when integrated, to provide a functional product, but are functional to effect integration.

Thus, the mtr gene sequences contained in the expression vector have at least two functions:

(1) they effect homologous integration of the inserted or contiguous DNA comprising the coding sequence for the desired protein, and

(2) they are of such a nature that when homologous combination occurs, no functional mtr protein is produced. Thus, they cannot themselves be sufficient to restore mtr protein function.

The construction of expression systems of the invention are illustrated by the initial construction of a host vector, designated herein pXpress, which includes the mtr gene sequence shown in

Figure 1. This vector is suitable for amplification in E. coli, and provides endogenous insertion sites for the DNA containing the desired coding sequence. Also illustrative of host vectors is pDXpress, an
5 analogous vector which is identical to pXpress except that an additional polylinker in the original host vector, from which pXpress was constructed, is deleted.

The illustrative vector pXpress is
10 constructed from the vector pBN3 which is described in Stuart, W.D., et al., Genome (1988) 30:198-203. pBN3 contains the 2783 bp N. crassa genomic DNA containing the mtr gene shown in Figure 1, which is bracketed by a BglII site in the gene and a BamHI site contained in
15 the vector. pBN3 was digested with BamHI and BglII and the segment containing the mtr gene was inserted into the BamHI site of the commercially available vector, pTZ18R, obtained from Pharmacia. This produced clones pN807 and pN816 wherein the EcoRI site
20 contained in the polylinker of the pTZ18R vector is upstream of the ORF; clones in the opposite orientation were designated pN846 and pN839. pXpress is a version of pN846 wherein the 5' polylinker of pTZ18R is deleted. The pXpress vector has useful
25 cloning sites for insertion of the desired DNA in the upstream region just downstream of the mtr promoter (SalI/AccI/HincII) (position 307 in Figure 1) and also in the latter third of the ORF (HincII at position 1406 and AccI at position 1920).

30 Thus, in one illustrative embodiment, pXpress may be used directly as an insertion vector to construct the expression system of the invention by cleaving with AccI, which deletes most of the open reading frame while retaining both the promoter and
35 transcription terminator of the mtr gene. In another embodiment, the vector may first be digested with SalI, which recognizes only the upstream post-promoter site, and then with HincII which, since the upstream recognition sites overlap, now recognizes only the

downstream site to produce a SalI-blunt opening into which the desired DNA may be inserted in a directionally controlled way.

pDXpress is a modified form of pXpress in which the linker provided in the host vector pTZ18R at the 3' end of the pN846 insert is deleted, thus eliminating additional restriction sites duplicate to those available in the sequence of the mtr gene.

After construction of a suitable expression vector by insertion of the desired coding sequence into the host vector containing the nucleotide sequence derived from mtr, cells of the desired recombinant host are treated with the vector under conditions which favor the uptake of the vector DNA into the cells. These transformation conditions are standard and depend on the nature of the cells to be transformed. For transformation of filamentous fungi, the preferred hosts of the invention, spheroplasts or protoplasts are generally used and the vector is linearized prior to treating the cells. The transformants are then selected on medium containing a substance to which mtr confers susceptibility, such as pFPA or 4-MT. Successful homokaryotic transformants can survive in this medium and can be cloned.

The cloned cells lines are then used in culture for production of recombinant protein encoded by the inserted DNA. The protein thus produced may be contained intracellularly or secreted into the medium, depending on the expression system construction used, as is understood in the art. After culturing under conditions which encourage the production of the protein, the protein produced is recovered from the culture and, if desired, purified using standard techniques known in the art. Alternatively, the cultures themselves or the culture media may provide the protein in sufficient purity for its intended use, as may be the case, for example, for the production of industrial enzymes or enzymes used for organic synthesis steps. Of course, if the protein produced

is intended for pharmaceutical use, it may be desirable to purify the protein produced to a suitable level for formulation.

The following examples illustrate, but do not limit the invention.

Example 1

Preparation of pBN3

11.3 μ g of BglII digested 5:4H DNA (DNA from cosmid 5:4H in the Vollmer-Yanosky library, described in Vollmer et al., PNAS (1986) 83:4869-4873), was electrophoresed in a preparative 1% LMT-agarose gel. 15.0 μ l (70 ng of DNA) of the liquified agarose gel, containing the 15.0 kbp fragment was mixed with 1.5 μ l H₂O, 2.0 μ l 10X T4 ligase buffer at 37°C and maintained at this temperature for 15 min. The cosmid 5:4H is capable of transforming an mtr negative host strain, as are 5:3B and 5:10H from the same library. Then 1.5 μ l 1.4 U/ μ l T4 DNA ligase was added. The contents were briefly mixed by flicking the tube and brought down with a quick pulse in a microcentrifuge. The reaction was incubated at 15°C overnight.

At the end of the incubation, 150 μ l Tris. Cl/CaCl₂ was added to the ligation reaction which was heated to 65-70°C for 5 minutes to liquify the agarose. The condensation was brought down with a rapid pulse, and the tube was immersed into an ice bath. The contents, after sufficient cooling, were dispensed into a microfuge tube containing 200 μ l competent E. coli LE392 cells. The transformation and plating of the cell/media mix was the same as previously described.

Example 2

Generation of pN807 and pN846

6 μ g pTZ18R DNA as digested to completion with 17 U BamHI and the final concentration adjusted to 60 ng/ μ l. 12.0 μ g pBN3 DNA was double digested with 36 U BglII and 34 U BamHI. The sample was

electrophoresed in a 1% LMT-agarose gel and the 2.9 kbp fragment was excised. The agarose plug was immersed in H₂O at room temperature (60 min.) to remove excess EtBr and 1X TBE buffer. The agarose plug, containing 600 ng of the 2.9 kbp fragment was liquified by heating and mixed with 120 ng of BamHI digested pTZ18R vector. 5X T4 ligase buffer and 2 U T4 DNA ligase was mixed into the solution at 37°C for a final volume of 90 µl. The reaction was incubated at room temperature (overnight).

A half volume aliquot of the reaction mix was liquified and mixed with an equal volume of Tris Cl/CaCl₂ solution, and cooled in an ice bath. The solution was used to transform 200 µl competent E. coli NM522 cells. The transformed cells were spread on X-gal indicator plates to detect recombinant DNA molecules. The numbered clones pN389 and pN846 are the same clones picked and numbered independently from different colonies. The numbered clones pN807 and pN816 are the same clones picked and numbered independently from different colonies, and contain the pBN3 insert in opposite orientation from that of pN839/846?

25

Example 3

Preparation of pXpress and pDXpress

Five micrograms of pN846 DNA were isolated from E. coli NM522 by standard methods (Koo and Stuart, 1991). The DNA was double digested with XbaI and HindIII, treated with Klenow and NTPs, cleaned with GeneClean (Bio 101), and ligated with 400 units of DNA T4 polymerase at room temperature overnight. The ligation mixture was used to transform E. coli NM522 host cells and selected for Amp^r. Transformed colonies were picked and grown in 1.5 ml liquid cultures in tubes overnight. Plasmid DNA was isolated and tested for the presence of HindIII, XbaI and PstI restriction sites. Isolates which had lost the three sites were then tested for the remaining sites

expected to be in pN846. One plasmid which had lost the expected sites and retained the expected sites was designated plasmid pXpress.

To prepare pDXpress, pXpress was digested
5 with BamHI and EcoRI, the ends polished with Klenow,
and the DNA is relegated (in the presence of SmaI, to
reduce the number of copies of partially cut and
relegated vector). The ligation mixture is
transformed into E. coli host cells and transformants
10 selected for resistance to ampicillin. Plasmid DNA
from resistant colonies is tested for loss of EcoRI,
BamHI and SacI sites. A plasmid meeting these
criteria and also retaining the internal KpnI sites of
pXpress was designated pDXpress (plasmid downstream
15 express).

Example 4

Production of Chymosin

The gene encoding bovine chymosin (Moir et
20 al., Gene (1982) 29:127-138) was inserted into pXpress
as a XhoI/HindIII-blunted fragment. pXpress was
digested with SalI and then with HincII, and blunted.
This deletes the portion of the mtr gene between
positions 307 and 1406 shown in Figure 1.

25 A sample of 0.5 μ g of the chymosin open
reading frame was ligated into 0.5 μ g of the cleaved
vector using 40 units of T4 ligase incubated overnight
at 16°C. Some fragments ligate SalI to XhoI and blunt
HindIII to blunt HincII (losing all four restriction
30 sites).

The ligated fragments were transformed into
competent E. coli cells DH-5alpha and selected for
Amp^r. Plasmids were isolated and digested with EcoRV
to confirm the insert. Plasmids containing the insert
35 in the correct orientation were renamed pNEC for
plasmid neutral (mtr) expression of chymosin.

Transformation of pNEC into Neurospora
spheroplasts was accomplished by standard methods.
See Koo and Stuart (1991) supra. pNEC DNA was

linearized by cutting with SacI and 5 μ g of linearized pNEC was used to transform 1×10^8 spheroplasts of strain 74a "wild" type Neurospora (Fungal Genetics Stock Center, Dept. of Microbiology, University of Kansas Medical Center, Kansas City, Kansas 66103).
5 The mixture was taken up in 15 ml of minimal top agar and spread onto a bottom plate containing 0.05 mg/ml pFPA. Plates were screened 3 days later. Eighteen colonies were picked and grown on solid Vogel's 1X
10 media containing 0.05 mg/ml pFPA in tube slants.

Colonies were transformed to liquid cultures of 1X Vogel's with 2% sucrose in double distilled H₂O. The liquid was collected and assayed for chymosin activity using 25 μ l culture, 125 μ l 10 mM Phosphate
15 Buffered Sodium (PBS), and 150 μ l clotting buffer (4.5 mg/ml CaCl₂, 3.0 mg/ml NaOAc in 10 mg/ml nonfat milk). Samples were tested on a 64 well microtiter plate, incubated at 37°C for 15 min then assayed by
absorbance at 630 nm on a Dynatech MR 5000 ELISA
20 machine. Among the 18 transformants, four showed milk clotting ability. Chymosin concentrations ranging from 1.6-4.6 μ g/ml were extrapolated from known chymosin standards.

ELISA assay was performed on the pNEC
25 transformants using rabbit antiovine chymosin antibody. Secondary antibody, goat antirabbit Ab conjugated to alkaline phosphatase was used to visualize chymosin concentrations. The ELISA reactions indicated chymosin concentrations ranging
30 from 1-16 μ g/ml. These results indicated that not all of the chymosin product was biologically active. Western blots of PAGE gels showed that some of the chymosin in the Neurospora media was in the form of prochymosin which is inactive until converted by
35 incubation at pH4 for several hours.

Example 5Expression of Relaxin Gene

A gene coding for the open reading frame of porcine relaxin (Haley et al., DNA (1982) 1:155-162) was inserted into the pXpress in a manner similar to the protocols used for chymosin. The gene was copied using the PCR reaction using primers designed so that a KpnI site and the Neurospora translation consensus sequence (Koo and Stuart 1991) were added to the 5' end of the gene and a BamHI site was added to the 3' end. The PCR product was polished with T4 polymerase and inserted in the plasmid pXpress at the HincII sites in a manner identical to the chymosin construction.

Orientation was confirmed by double digestion with EcoRV and BamHI and with single digestion with KpnI. Inserts with 5' end of the gene following the mtr promoter were used to transform Neurospora spheroplasts as in Example 4.

Transformants resistant to pFPA were transferred to liquid cultures and the media screened for production of relaxin by ELISA assay using a rabbit anti-relaxin antibody and goat anti-rabbit secondary antibody. Six transformants secreted a protein which cross reacted with relaxin antibody, but at levels 10 to 100 times lower than chymosin as determined by ELISA.

30

35

CLAIMS

1. An expression system for expressing a gene encoding a desired protein in a recombinant host having a functional mtr locus, said expression system consisting of a DNA molecule which molecule comprises:
- 5 a first nucleotide sequence encoding said desired protein;
- a second nucleotide sequence derived from
- 10 the mtr gene of N. crassa disposed in relationship to said first sequence so as to permit homologous recombination of said first and second sequences into the mtr locus of said host so as to incapacitate the mtr locus in said host from effecting production of
- 15 functional mtr protein;
- wherein said first nucleotide sequence is operably linked to control sequences which effect expression of said encoding sequence, when contained in the recombinant host cell.
- 20
2. The expression system of claim 1 wherein at least a portion of said second sequence is disposed at the 5' terminus of the said first nucleotide sequence.
- 25
3. The expression system of claim 2 wherein said portion comprises the mtr promoter operably linked to said first nucleotide sequence.
- 30
4. The expression system of claim 1 wherein at least a portion of said second nucleotide sequence is disposed at the 3' terminus of said first nucleotide sequence.
- 35
5. The expression system of claim 4 wherein said portion comprises the mtr transcription terminator sequences operably linked to said first nucleotide sequence.

6. The expression system of claim 1 wherein said first nucleotide sequence is operably linked to a promoter homologous to said first nucleotide sequence, or wherein said first DNA
5 sequence is operably linked to a promoter which is heterologous both to said first and second nucleotide sequences, or

wherein said first nucleotide sequence is operably linked to a transcription terminator
10 homologous to said first nucleotide sequence, or wherein said first DNA sequence is operably linked to a transcription terminator heterologous to both said first and second nucleotide sequences, or wherein said first nucleotide sequence is an
15 open reading frame with the coding region of said second sequence, or

wherein said second nucleotide sequence comprises less than the complete sequence of the mtr gene.

20

7. A recombinant host cell which contains the expression system of claim 1.

8. The cell of claim 7 which is Neurospora
25 crassa.

9. A method to produce a desired protein which method comprises culturing the host cell of claim 7 under conditions wherein expression of the
30 nucleotide sequence encoding the desired protein is effected to as to produce said protein in the culture; and

recovering the protein from the culture.

35 10. The method of claim 9 wherein said desired protein is relaxin or chymosin.

11. A method to select for a recombinant host cell which has, integrated into all mtr loci in

said cell, the expression system of claim 1, which method comprises culturing said candidate cells on a medium containing a substance which is lethal to said cell in the presence of a functional mtr gene product;

5 and

selecting cells capable of growth on said medium.

12. A method to obtain cells which are
10 homokaryotic transformants with a desired DNA, which method comprises contacting said host cells,
in a form susceptible for uptake of DNA molecules and under conditions which are suitable for the uptake of DNA molecules,

15 with a DNA molecule which comprises said desired DNA contiguous to a sequence homologous to a recessive gene that confers susceptibility to a poison, so as to permit homologous recombination of said DNA molecule into the locus of said recessive
20 gene to prevent expression of said gene;

culturing the resulting cells in a medium containing said poison; and

recovering cells which are capable of growth in the presence of said poison.

25

13. A method to obtain cells which are homokaryotic transformants with a desired DNA, which method comprises

culturing cells that have been treated, in a
30 form susceptible for uptake of DNA and under conditions which are suitable for the uptake of DNA molecules with a DNA molecule which comprises said desired DNA contiguous to a sequence homologous to a recessive gene that confers susceptibility to a
35 poison, so as to permit homologous recombination of said DNA molecule into the locus of said recessive gene to prevent expression of said gene;

in a medium containing said poison; and

recovering cells which are capable of growth
in the presence of said poison.

5

10

15

20

25

30

35

1/4

DNA Strider 1.0 ### Tuesday, June 15, 1993 10:07:28 AM

New 2.9 kb DNA seqn -> List

DNA sequence 2783 b.p. AGATCCGCCTCG ... GATGTTCCAGGC linear

1	AGATCCGCCT	10	CGCCCCAAGC	20	GCATCCCAAC	30	GCAGGCGTGC	40	TATATGTCGC	50	TCCTCCCTCT	60	TCCTCCCTCT
61	CACGTACCTC	70	GCAAGTACCT	80	GTCTCATCTG	90	GCACCCGCTT	100	TCCTCCATCCC	110	TCCTCCATCCC	120	TCCTCCATCCC
121	TACTTAACT	130	GCCCGGCTCC	140	CCCGCAGTTC	150	ATCCGCTCTC	160	TCAGACCTTG	170	GATCGTCTGT	180	GATCGTCTGT
181	TGGTCTTTTC	190	GCTTATTTTC	200	CGTACCTTTC	210	TCCTATTTTC	220	TCCTATTTTC	230	TCCTATTTTC	240	TCCTATTTTC
241	CAGCATTTTC	250	TCCTATTTTC	260	TCCTATTTTC	270	TCCTATTTTC	280	TCCTATTTTC	290	TCCTATTTTC	300	TCCTATTTTC
301	GCCACTGTTC	310	ACAAAGGCTC	320	TCAACAAGGC	330	CTCTTTACCA	340	CCCAACATGG	350	ACTCGCAATA	360	ACTCGCAATA
361	CGAGACAAAA	370	AAGAATGACC	380	CAAAAGGCTC	390	CATGCGGTAC	400	CCAGAGTCAA	410	ACGATGAGCA	420	ACGATGAGCA
421	TGTTGGCGAG	430	GTCCGCGGCT	440	TGGGCGGCTC	450	CATCATGGAC	460	AAGGAGCTTG	470	AGGCCCAAGG	480	AGGCCCAAGG
481	GGGCCAATGCC	490	AAGTTCACCC	500	GTCTCGGCTG	510	GAAACGCTTG	520	ACGGTCTGTC	530	TCATCGTCTG	540	TCATCGTCTG
541	GGCCATTTGCC	550	CTCGGCTCTC	560	TCTGCTTTCC	570	CGGCGCTTTC	580	GCTACCCCTTG	590	GCTACCCCTTG	600	GCTACCCCTTG
601	TGGTGTTATT	610	CTCTCTGTTC	620	GCAATGGGACT	630	CATCTGCATC	640	TACACGGGCTC	650	ACGTTATCGG	660	ACGTTATCGG
661	ACAAACCAAG	670	CTCAAGCACC	680	CTGAAATCGC	690	CCACTATGCC	700	GACGTTGGTC	710	GTGTCTATGT	720	GTGTCTATGT
721	TGGAAGATGG	730	GGATATGAAA	740	TCATCAGCTT	750	CATGTTTGTG	760	CTGCAACTGA	770	TCCTCATCGT	780	TCCTCATCGT
781	CGGCTCCAC	790	GTCCCTCACG	800	GCACCAATCAT	810	GTGGGGCACC	820	ATCACGGATA	830	ACGGCAACGG	840	ACGGCAACGG
841	TACCTGCTCT	850	CTCGTCTTTC	860	GCAATGCTCT	870	CGCCATCATT	880	CTCTTCTCTC	890	TTGCCATTCC	900	TTGCCATTCC
901	TCCCAGTTTC	910	GCCGAGTTTG	920	CCATCTTTGG	930	ATACATCGAT	940	TTGCTCTCCA	950	TCGTGCGCCGC	960	TCGTGCGCCGC
961	CATCTCATTC	970	ACCATGATTG	980	CTACTGGCAT	990	TCGCTCGAGC	1000	CACCAAGGAGG	1010	GTGGTCTCGC	1020	GTGGTCTCGC
1021	TGCTGTTCCC	1030	TGGTCTTTGCT	1040	GGCCCCAAGGA	1050	GGACCTTAGC	1060	CTTGCTGAGG	1070	GCTTTCATTGC	1080	GCTTTCATTGC

FIG.1A

SUBSTITUTE SHEET

2/4

1081	TGTCAGCAAC	ATCGTTTTTCG	CCTACAGCIT	CCCCATGTCC	CAGTTCAGCT	TTATGGATGA	1140
1141	GATGCACACC	CCCTCCGACT	ACAAGAAAGIC	CATCGTTGCT	CTCGGCTTGA	TTGAAATCTT	1200
1201	CATCTACACC	GTTACTGGG	GCCTCGTTTA	CGCTTTCGTC	GGCCCCGAGG	TCCAGTCTCC	1260
1261	TGCTTGTCTC	TCTGCTGGCC	CTCTCTCTGC	CAAGGTGCTC	TTGCGCATTG	CCCTCCCCGT	1320
1321	CATCTTCAIC	TCTGGCAGTA	TCAACACIGT	TGCTGTCAGC	AGGTATCTGA	TTGAGCGCAT	1380
1381	CTGGCCCAAC	AACGTCAATC	GCTATGTCAA	CACCCACGCG	GGTTGGATGG	TTTGGCTTGG	1440
1441	TTTTGACTTT	GGCATTAACCC	TCAATGCCIG	GGTATATGCT	GAGGCCATCC	CTTTCTTCTC	1500
1501	TGATCTGTTG	GCCATCTGCT	CGGCTCTCTT	CATTTCCGGT	TTTAGCTTCT	ATTTCCCTGC	1560
1561	CTTGATGTAT	TTCAAGATCA	CCAGGAACGA	TCCCAAGAGC	CAGGGCAACA	AGTACTTCTT	1620
1621	GGATGCCCTC	AACATGCTCT	GCTTCGTCT	CGGCATGGGC	ATTTCTGGTA	TTGGTACCTA	1680
1681	CGCCGCTATT	CAGGACATTG	TAAGTTTGGC	CCGCTTTTCT	GTTTACTCTT	TGCACACAAA	1740
1741	TGCTAACTTG	CTTCTCAGAT	GGACCGTTAC	GACCAATGGCA	AGGTTTCGAA	GCCTTATAGC	1800
1801	TGTGCGCCCT	TGGCTTAACA	GGCCCAACGC	ACGCTTATGA	TCCCTGTTGT	TTTTTTTGGG	1860
1861	TGATTTAATT	AAAGTTGCGC	AGTGATTGAC	GTCGTCTTTC	ACCCGCGATT	TTTTTCGGAG	1920
1921	TATACCCCTT	CAGACTTGCC	GGCTTGGGGA	AATGTTTTGA	GTTATTTCTAT	TTTCGGAGTT	1980
1981	TCAGGATTTG	GCACAAAGCA	AACCAAGCGG	GAGTTGAAAC	CGTGGTGTGG	TCGCGGTGCG	2040
2041	CTGCTGCATT	GGTAGTGCTT	GTTCAGGTTT	TTTGGTTGGT	GGTTTGGATG	CGTGACCCAC	2100
2101	TTTTTTTTTT	AACGTTTTAT	TGCATGCATG	TATTATATGG	GAAAGTCATG	GGACATGGCA	2160
2161	ACTATACGAA	CCGACGCAAA	GATAGGATGG	GATGGATGAT	GGATGGACGT	ACGATCCAAC	2220
2221	CGCTGGGGA	CTGGACTGAA	CGGAATTAGG	ACGGACGGGA	CAGGTAACCT	AGGTACCTAA	2280
2281	TGACCGGAAT	ATGTTTACAA	ATCATTTGTT	AGTGCGGGTG	ACCGGCAATA	GAGACGATGG	2340
2341	GCACAGGAAT	ATCGATAGAT	GCTACCTATA	CTCTAAAGAA	CTCTATAGGT	ATAATAATCG	2400
2401	CTGAACATAC	CTTGCCCCAA	TTTCCATCCT	ACACCCATGG	TTATGAATC	ATCCIGTTGT	2460
2461	TGTGCCATAA	TTTTCCATCT	GACTCCCAATG	CTTCTTCTCT	TTTTTTTTTT	TTTTTCTTGT	2520
2521	GCCACGCGCC	TCCGATATCT	CGAGTTTTTG	AAGGATTTCT	GIGTTGGGTG	GAGCTTTTCT	2580
2581	CAACAAATCCT	GGGGCTTCGA	ATCCCTCCAC	CAGACCTCAC	CCCAGCAGTC	AGAGTTTAGC	2640
2641	CCGCCAGCCA	GCCCGTTCAGC	CAAGTGAGTT	CTAAGATTAA	TCTCGACTCC	TTGACAAAGGC	2700
2701	TTTGCGTGGC	CACGCTCTCCT	CTCAACACGC	AAAACTTTTG	TCATTGTTAC	TACACTACAG	2760
2761	GTTACCGTGT	CGATGTTCCA	GGC				2783

FIG.1B

SUBSTITUTE SHEET

3/4

FIG.2A

New 2.9 kb DNA seqn [347 to 1699] → 1-phase Translation

DNA sequence 2783 b.p. AGATCCGGCTCG ... GATGTTCCAGGC linear

347	ATG	met	407	TCA	ser	467	CCT	pro	527	GTC	val	587	CTT	leu	647	GCT	ala	707	GGT	gly	767	CTG	leu	827	GAT	asp	887	CTC	leu	947	TCC	ser	1007	GAG	glu	1067	GAG	glu																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
1	TCG	ser	21	AAC	asn	41	GAG	glu	61	CTC	leu	81	GGC	gly	101	CAC	his	121	CGT	arg	141	ATC	leu	161	AAC	asn	181	CTT	leu	201	ATC	ile	221	GGT	gly	241	GGC	gly																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
377	/	11	377	/	11	437	/	31	497	/	51	557	/	71	617	/	91	737	/	131	857	/	171	917	/	211	1037	/	231	1097	/	251	1157	/	1171	1217	/	1231	1297	/	1311	1377	/	1391	1457	/	1471	1537	/	1551	1617	/	1631	1697	/	1711	1777	/	1791	1857	/	1871	1937	/	1951	2017	/	2031	2097	/	2111	2177	/	2191	2257	/	2271	2337	/	2351	2417	/	2431	2497	/	2511	2577	/	2591	2657	/	2671	2737	/	2751	2817	/	2831	2897	/	2911	2977	/	2991	3057	/	3071	3137	/	3151	3217	/	3231	3297	/	3311	3377	/	3391	3457	/	3471	3537	/	3551	3617	/	3631	3697	/	3711	3777	/	3791	3857	/	3871	3937	/	3951	4017	/	4031	4097	/	4111	4177	/	4191	4257	/	4271	4337	/	4351	4417	/	4431	4497	/	4511	4577	/	4591	4657	/	4671	4737	/	4751	4817	/	4831	4897	/	4911	4977	/	4991	5057	/	5071	5137	/	5151	5217	/	5231	5297	/	5251	5317	/	5331	5397	/	5351	5417	/	5431	5497	/	5451	5517	/	5531	5597	/	5551	5617	/	5631	5697	/	5651	5717	/	5731	5797	/	5751	5817	/	5831	5897	/	5911	5977	/	5991	6057	/	6071	6137	/	6151	6217	/	6231	6297	/	6251	6317	/	6331	6397	/	6351	6417	/	6431	6497	/	6451	6517	/	6531	6597	/	6551	6617	/	6631	6697	/	6651	6717	/	6731	6797	/	6751	6817	/	6831	6897	/	6851	6917	/	6931	6997	/	7011	7077	/	7031	7097	/	7051	7117	/	7131	7197	/	7151	7217	/	7231	7297	/	7251	7317	/	7331	7397	/	7351	7417	/	7431	7497	/	7451	7517	/	7531	7597	/	7551	7617	/	7631	7697	/	7651	7717	/	7731	7797	/	7751	7817	/	7831	7897	/	7851	7917	/	7931	7997	/	8011	8077	/	8031	8097	/	8051	8117	/	8131	8197	/	8151	8217	/	8231	8297	/	8251	8317	/	8331	8397	/	8351	8417	/	8431	8497	/	8451	8517	/	8531	8597	/	8551	8617	/	8631	8697	/	8651	8717	/	8731	8797	/	8751	8817	/	8831	8897	/	8851	8917	/	8931	8997	/	8951	9017	/	9031	9097	/	9051	9117	/	9131	9197	/	9151	9217	/	9231	9297	/	9251	9317	/	9331	9397	/	9351	9417	/	9431	9497	/	9451	9517	/	9531	9597	/	9551	9617	/	9631	9697	/	9651	9717	/	9731	9797	/	9751	9817	/	9831	9897	/	9851	9917	/	9931	9997	/	10011	10077	/	10031	10097	/	10051	10117	/	10131	10197	/	10151	10217	/	10231	10297	/	10251	10317	/	10331	10397	/	10351	10417	/	10431	10497	/	10451	10517	/	10531	10597	/	10551	10617	/	10631	10697	/	10651	10717	/	10731	10797	/	10751	10817	/	10831	10897	/	10851	10917	/	10931	10997	/	10951	11017	/	11031	11097	/	11051	11117	/	11131	11197	/	11151	11217	/	11231	11297	/	11251	11317	/	11331	11397	/	11351	11417	/	11431	11497	/	11451	11517	/	11531	11597	/	11551	11617	/	11631	11697	/	11651	11717	/	11731	11797	/	11751	11817	/	11831	11897	/	11851	11917	/	11931	11997	/	11951	12017	/	12031	12097	/	12051	12117	/	12131	12197	/	12151	12217	/	12231	12297	/	12251	12317	/	12331	12397	/	12351	12417	/	12431	12497	/	12451	12517	/	12531	12597	/	12551	12617	/	12631	12697	/	12651	12717	/	12731	12797	/	12751	12817	/	12831	12897	/	12851	12917	/	12931	12997	/	12951	13017	/	13031	13097	/	13051	13117	/	13131	13197	/	13151	13217	/	13231	13297	/	13251	13317	/	13331	13397	/	13351	13417	/	13431	13497	/	13451	13517	/	13531	13597	/	13551	13617	/	13631	13697	/	13651	13717	/	13731	13797	/	13751	13817	/	13831	13897	/	13851	13917	/	13931	13997	/	13951	14017	/	14031	14097	/	14051	14117	/	14131	14197	/	14151	14217	/	14231	14297	/	14251	14317	/	14331	14397	/	14351	14417	/	14431	14497	/	14451	14517	/	14531	14597	/	14551	14617	/	14631	14697	/	14651	14717	/	14731	14797	/	14751	14817	/	14831	14897	/	14851	14917	/	14931	14997	/	14951	15017	/	15031	15097	/	15051	15117	/	15131	15197	/	15151	15217	/	15231	15297	/	15251	15317	/	15331	15397	/	15351	15417	/	15431	15497	/	15451	15517	/	15531	15597	/	15551	15617	/	15631	15697	/	15651	15717	/	15731	15797	/	15751	15817	/	15831	15897	/	15851	15917	/	15931	15997	/	15951	16017	/	16031	16097	/	16051	16117	/	16131	16197	/	16151	16217	/	16231	16297	/	16251	16317	/	16331	16397	/	16351	16417	/	16431	16497	/	16451	16517	/	16531	16597	/	16551	16617	/	16631	16697	/	16651	16717	/	16731	16797	/	16751	16817	/	16831	16897	/	16851	16917	/	16931	16997	/	16951	17017	/	17031	17097	/	17051	17117	/	17131	17197	/	17151	17217	/	17231	17297	/	17251	17317	/	17331	17397	/	17351	17417	/	17431	17497	/	17451	17517	/	17531	17597	/	17551	17617	/	17631	17697	/	17651	17717	/	17731	17797	/	17751	17817	/	17831	17897	/	17851	17917	/	17931	17997	/	17951	18017	/	18031	18097	/	18051	18117	/	18131	18197	/	18151	18217	/	18231	18297	/	18251	18317	/	18331	18397	/	18351	18417	/	18431	18497	/	18451	18517	/	18531	18597	/	18551	18617	/	18631	18697	/	18651	18717	/	18731	18797	/	18751	18817	/	18831	18897	/	18851	18917	/	18931	18997	/	18951	19017	/	19031	19097	/	19051	19117	/	19131	19197	/	19151	19217	/	19231	19297	/	19251	19317	/	19331	19397	/	19351	19417	/	19431	19497	/	19451	19517	/	19531	19597	/	19551	19617	/	19631	19697	/	19651	19717	/	19731	19797	/	19751	19817	/	19831	19897	/	19851	19917	/	19931	19997	/	19951	20017	/	20031	20097	/	20051	20117	/	20131	20197	/	20151	20217	/	20231	20297	/	20251	20317	/	20331	20397	/	20351	20417	/	20431	20497	/	20451	20517	/	20531	20597	/	20551	20617	/	20631	20697	/	20651	20717	/	20731	20797	/	20751	20817	/	20831	20897	/	20851	20917	/	20931	20997	/	20951	21017	/	21031	21097	/	21051	21117	/	21131	21197	/	21151	21217	/	21231	21297	/	21251	21317	/	21331	21397	/	21351	21417	/	21431	21497	/	21451	21517	/	21531	21597	/	21551	21617	/	21631	21697	/	21651	21717	/	21731	21797	/	21751	21817	/	21831	21897	/	21851	21917	/	21931	21997	/	21951	22017	/	22031	22097	/	22051	22117	/	22131	22197	/	22151	22217	/	22231	22297	/	22251	22317	/	22331	22397	/	22351	22417	/	22431	22497	/	22451	22517	/	22531	22597	/	22551	22617	/	22631	22697	/	22651	22717	/	22731	22797	/	22751	22817	/	22831	22897	/	22851	22917	/	22931	22997	/	22951	23017	/	23031	23097	/	23051	23117	/	23131	23197	/	23151	23217	/	23231	23297	/	23251	23317	/	23331	23397	/	23351	23417	/	23431	23497	/	23451	23517	/	23531	23597	/	23551	23617	/	23631	23697	/	23651	23717	/	23731	23797	/	23751	23817	/	23831	23897	/	23851	23917	/	23931	23997	/	23951	24017	/	24031	24097	/	24051	24117	/	24131	24197	/	24151	24217	/	24231	24297	/	24251	24317	/	24331	24397	/	24351	24417	/	24431	24497	/	24451	24517	/	24531	24597	/	24551	24617	/	24631	24697	/	24651	24717	/	24731	24797	/	24751	24817	/	24831	24897	/	24851	24917	/	24931	24997	/	24951	25017	/	25031	25097	/	25051	25117	/	25131	25197	/	25151	25217	/	25231	25297	/	25251	25317	/	25331	25397	/	25351	25417	/	25431	25497	/	25451	25517	/	25531	25597	/	25551	25617	/	25631	25697	/	25651	25717	/	25731	25797	/	25751	25817	/	25831	25897	/	25851	25917	/	25931	25997	/	25

4/4

1127 / 261
 AGC TTT ATG
 ser phe met
 1187 / 281
 TTG ATT GAA
 leu ile glu
 1247 / 301
 GAG GTC CAG
 glu val gln
 1307 / 321
 ATT GCC CTC
 ile ala leu
 1367 / 341
 CTG ATT GAG
 leu ile glu
 1427 / 361
 ATG GTT TGG
 met val trp
 1487 / 381
 ATC CCT TTC
 ile pro phe
 1547 / 401
 TTC TAT TTC
 phe tyr phe
 1607 / 421
 AAG AAG TAC
 lys lys tyr
 1667 / 441
 GGT ATT GGT
 gly ile gly
 1727 /
 TCT TTG CAC
 GAT GAG ATG CAC ACC CCC TCC
 asp glu met his thr pro ser
 ATC TTC ATC TAC ACC GTT ACT
 ile phe ile tyr thr val thr
 TCT CCT GCC TTG CTC TCT GCT
 ser pro ala leu leu ser ala
 CCC GTC ATC TTC ATC TCT GGC
 pro val ile phe ile ser gly
 CGC ATC TGG CCC AAC AAC GTC
 arg ile trp pro asn val
 CTT GGT TTT GAC TTT GGC ATT
 leu gly phe asp phe gly ile
 TTC TCT GAT CTG TTG GCC ATC
 phe ser asp leu leu ala ile
 CCT GCC TTG ATG TAT TTC AAG
 pro ala leu met tyr phe lys
 TTC TTG GAT GCC CTC AAC ATG
 phe leu asp ala leu asn met
 ACC TAC GCC GCT ATT CAG GAC
 thr tyr ala ala ile gln asp
 1157 / 271
 GAC TAC AAG
 asp tyr lys
 1217 / 291
 GGT GGC GTC
 gly gly val
 1277 / 311
 GGC CCT CTT
 gly pro leu
 1337 / 331
 AGT ATC AAC
 ser ile asn
 1397 / 351
 ATT CGC TAT
 ile arg tyr
 1457 / 371
 ACC CTC ATT
 thr leu ile
 1517 / 391
 TGC TCG GCT
 cys ser ala
 1577 / 411
 ATC ACC AGG
 ile thr arg
 1637 / 431
 CTC TGC TTC
 leu cys phe
 1697 / 451
 ATT GTA AGT
 ile
 1759 /
 ATG GAC CAT GGC AAG GTT
 met asp arg tyr asp his gly lys val

FIG.2B

1789 /
 TCC AAG CCT TAT AGC TGT GCG CCC TTG GCT TAA
 ser lys pro tyr ser cys ala pro leu ala och

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/05866

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C12N 5/10, 15/16, 15/80

US CL : 435/69.1, 69.4, 172.3, 254.11, 254.4, 320.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 69.4, 172.3, 254.11, 254.4, 320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Dialog (Biosis, Medline, CAS, Biotech Abstr.)

APS

Search terms: MTR and Neurospora

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Genome Vol. 30, No. 2, issued 1988, Stuart et al., "Cloning of mtr an amino acid transport gene of <u>Neurospora crassa</u> ", pages 198-203, see abstract no. 6581989.	1-13
Y	Genetics Vol. 116 (1 part 2), issued 1987, Koo et al., "Cloning of the mtr gene", page 531, see abstract no. 7.6.	1-13
Y	Genome Vol. 34, No. 4, issued 1991, Koo et al., "Sequence and structure of mtr an amino acid transport gene of <u>Neurospora crassa</u> ", pages 644-651, see abstract no. 8671122.	1-13

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	* T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A* document defining the general state of the art which is not considered to be part of particular relevance	* X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E* earlier document published on or after the international filing date	* Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* &*	document member of the same patent family
* O* document referring to an oral disclosure, use, exhibition or other means		
* P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 04 August 1993	Date of mailing of the international search report 26 AUG 1993
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer JOHN LEGUAYDER
Facsimile No. NOT APPLICABLE	Telephone No. (703) 308-0196